

Some redox titration problems involving manganate(VII)

Background skills

1. How many moles of manganate(VII) ions, MnO_4^- , are in the following solutions?

- a) 1000cm^3 of 1M potassium manganate(VII), KMnO_4 .
- b) 25.0cm^3 of 1M KMnO_4 .
- c) 25.0cm^3 of 0.020M KMnO_4 .
- d) 36.5cm^3 of 0.012M KMnO_4 .

2. Calculate the relative formula masses of the following substances using A_r values from the table on the right.

- a) MnO_4^- ion
- b) KMnO_4

Element	A_r
H	1.00
N	14.0
O	16.0
S	32.1
K	39.1
Mn	54.9
Fe	55.8

3. What are the concentrations of MnO_4^- ions in the following solutions?

- a) 1 mole of KMnO_4 dissolved in 1000cm^3 of water.
- b) 0.05 moles of KMnO_4^- ions dissolved in 25cm^3 water.
- c) 39.5g of KMnO_4 dissolved in 250cm^3 water.
- d) 0.253g of KMnO_4 dissolved in 25.3cm^3 water.

4. a) Combine the following two half-reaction equations to give a balanced redox equation:



b) How many moles of Fe^{2+} ions are oxidised by 1 mole of MnO_4^- ions?

Titration problems

5. 25.0cm^3 of an acidified solution containing Fe^{2+} ions was titrated against potassium manganate(VII) solution. 20.0cm^3 of 0.050M potassium manganate(VII) was needed.

Calculate the concentration of Fe^{2+} ions in the acidified solution.

6. A 25.0cm^3 aliquot of a solution containing Fe^{2+} ions and Fe^{3+} ions was acidified and titrated against potassium manganate(VII) solution. 15.0cm^3 of 0.020M potassium manganate(VII) was needed. A second 25.0cm^3 aliquot was reduced using zinc (i.e. the Fe^{3+} ions in the solution were reduced to Fe^{2+} ions), then titrated. This time, 19.0cm^3 of the 0.020M potassium manganate(VII) was needed. Calculate the concentrations of:

- a) Fe^{2+} ions in the solution
- b) Fe^{3+} ions in the solution

7. *Kilzemaal* (a new fertiliser) contains ammonium iron(II) sulphate, $\text{FeSO}_4 \cdot (\text{NH}_4)_2\text{SO}_4 \cdot 6\text{H}_2\text{O}$ as a source of iron. A 6.50g sample of *Kilzemaal* is made up to 250cm^3 with dilute sulphuric acid. 25cm^3 of this solution reacted with 23.5cm^3 of 0.010M potassium manganate(VII). Calculate:

- a) The concentration of Fe^{2+} ions in the 25cm^3 aliquot.
- b) The number of moles of Fe^{2+} in the original 6.50g sample of *Kilzemaal*.
- c) If it contains more than 10.00% of iron by mass, *Kilzemaal* will kill 'em all. Will it?